

DETECTABLE SYSTEM OF RESISTANCE OF VERTIMEC (ABAMECTIN) WITHIN THE TWO-SPOTTED SPIDER MITE, *TETRANYCHUS URTICAE* KOCH

Rania Ahmed Abd El-Wahab
Plant Protection Research Institute, Agricultural Research Center, EGYPT
rania-proline@hotmail.com

Abstract - Augmented resistance of pesticides remains a principal issue for unsatisfactory impacts on the environment. So, there is a need to distinguish it faster by innovated devices and then find effective tools to reduce it as soon as possible, especially for mites such as *Tetranychus urticae*. Detectable tools, which are powered with solar energy, were used to discover Vertimec resistance depending on the assurance of the changes of electromagnetic field forces in comparison with sensitive mites. Acquired results with certain differences of electromagnetic forces were affirmed by evaluation of resistance proportions (RRs) which recorded 8.52- and 4.53-folds for resistant strains of Vertimec (Abamectin) on cotton and soybean, respectively, upon Vertimec LC50s. Furthermore, they were 5.97- and 4.29- folds for field Vertimec strains on cotton and soybean, respectively. However, there were no significant differences between resistant and field strains; they appeared in comparing the mentioned strains with susceptible mites. It may be well concluded that pesticides' resistance is now available to be detected by utilizing a simple and quick framework in the field.

Keywords - Detector, Electromagnetic, *Tetranychus urticae*, Resistance and Vertimec.

I. INTRODUCTION

Resistance of pesticides is an essential subject particularly when it is related to the two spotted spider mite, *Tetranychus urticae*. It is vital to take the most important components of climate changes as top priority when resistance is the issue which agriculturists should give a hand to. In Egypt, with elevated ratios of CO₂ and UV pro rata with pesticides' resistance specifically of spider mites, it is difficult to control them with pesticides (Nahar *et al.* 2005) due to their short life cycle, reproduction high capacity, and ability to create protection from miticides (Georghiou 1990).

Even with the broad utilization of chemical pesticides, pests existed a noteworthy explanation behind crop production calamity worldwide. Common used pesticides caused numerous ecological, agricultural, medical, and financial problems. Imperatively, development of resistance occurred as a result of the overwhelming utilization of synthetic insecticides and the selection pressure on insects and mites.

Particular metabolic pathways utilized by insects to change over hindrances into less poisonous structures or their expulsion from the framework are featured. Utilizing the proteomics approach, responsible proteins affected by pesticides in insects and then their alterations by pesticides' resistance could be distinguished (Dawkar *et al.*, 2013).

The available tool to assay pesticides' resistance is depending on Immunochromatographic dip-stick-format kits. They were created to distinguish resistance to carbamates and organophosphates and others. Strips were upon polyclonal antisera versus resistance of pesticides which are related to esterase isozymes isolated from insects. They were simply used by farmers anywhere (Kranthi, 2005), but the problem which is related to cost and accuracy still exists.

Using electromagnetic fields as traps or even to attract or eject pests, is an ordinary known process. It was started by Gerharz (1991) who situated the electromagnetic field nearby or in a pest control. A gadget was chosen from the gathering comprising insect traps, insect teasing stations containing a pest poison, and pointer stations. For instance, a wide assortment of regular pest traps for slithering insects could be retrofitted with an electromagnetic field generator. Devices could have the generator to a focal area of a trap with an electromagnetic field generator, contained or not, a chemical or biological attractant.

So, there is a need to get a priceless device that would be able to get fast and accurate results to help farmers to discover the problem in the beginning before its development. In the same time, there will be a new use of electromagnetic fields to detect pesticides' resistance in order to protect the environment of more quantities of acaricides.

Mites with certain changes in their bodies provided electric changes and then with the present magnetism in the instrument, electromagnetics field are available to react effectively with exposed mites. Nevertheless, resulted electromagnetic fields' forces and their Axis Graph Magnetic Meter were assessed by the innovated instrument in order to detect pesticides' resistance easily and vastly as presented in this paper.

II. MATERIALS AND METHODS

• Innovated Instrument Components:

The prototype of the instrument is shown in Image (1) with all of its components: A solar panel with its attachments and a DC-Motor. Electromagnetic field resulted from two magnets and with the passage of electric current, the required field was gained. Whenever the current is gone through mites, the electrophysiological differences appeared in changes of voltages at LCD monitor. Microcontroller system (PIC16F627A) was used to detect the resistance through electrochemical changes in the resistant strain in comparison with the other sensitive one. The sensor, UGN3503U was used to determine the magnetic field strength and varying voltage provided at output proportions which are picked up to the field strength. Also, Electromagnetic interference (EMI) detector was attached to the present circuit to provide and confirm accurate readings of the final electromagnetic field strength. Comparison appeared in the magnetic field force and the current conductance with attention to any differences in food resource, weather conditions, etc.

• Maintenance of *Tetranychus urticae* strains:

1. Susceptible strain

Colonies of the spider mite, *T.urticae* were raised under lab conditions ($25\pm 2^{\circ}\text{C}$, $60\pm 5\%\text{RH}$ and 12 hours light/12 hours dark) at Plant Protection Research Institute for several years without exposure to any contaminations or pesticides.

Image (1) Instrument

Prototype Parts:

- 1-Solar Panel
- 2-Container of battery, microcontroller system (PIC16F627A), UGN3503U, EMI and others
- 3-Power
- 4- Blue Light Emitting Diodes (LEDs)
- 5- Red Light Emitting Diodes (LEDs)



2. Resistant strain

Original colony of the spider mite, *T.urticae* was set up from mites gathered from castor oil plants without exposing to pesticides. It was raised under laboratory conditions ($25\pm 2^{\circ}\text{C}$, $60\pm 5\%\text{RH}$ and 12 hours light/12 hours dark) to assess the action of Vertimec (Abamectin $18\text{ g L}^{-1}\text{ EC}$) against *T.urticae* grown-up females. The leaf-dip technique presented by Dittrich (1962) was utilized. All treatments were done under laboratory conditions and each was replicated thrice. Likewise, control discs were dipped in water only. Mortality percentages were determined and corrected by using Abbott's formula (1925). Pooled data were subjected to probit analysis (POLO PC) (LeOra software, 1994). The original strain females were selected for Vertimec for 20 generations according to Yang et al. (2002) with some modifications. 1000 adult females of this colony started this selection. Every two generations, LC50s and LC90s were evaluated. New LC50 was applied as subsequent selection pressure. The next selection transferred to untreated leaves. LC50 estimations of the selected strain were compared to those of the susceptible strain. LC50 of Field colony was got after exposure to the recommended concentrations of Vertimec under certain values of UV and CO₂ which interacted with sprayed pesticide on infested cotton and soybean with *T.urticae*. Then, the resistance ratio (RR) was computed. Field colony was got after exposure to the recommended concentrations of Vertimec under certain values of UV and CO₂ which interacted with sprayed pesticide on infested cotton and soybean with *T.urticae*.

• **Determination of electromagnetic fields:**

Samples of *T.urticae*, about 100 adult females, were placed inside the instrument. Resistant and field strains were compared with the control which was reared on cotton and soybean leaves. Differences of electromagnetic fields' forces and Axis Graphs appeared and were recorded by the microcontroller system.

• **Data Analysis**

SPSS (V.16) was used to show differences among resistant, field and susceptible strains under electromagnetic' forces instrument. Jonckheere-Terpstra Test, Friedman Test, Kendall's W and others were used to test significance between resistance and susceptible cases at probability with 5% and 1%.

III. RESULTS AND DISCUSSION

Data revealed that strength of electromagnetic fields was changed particularly with significant differences between resistant and susceptible mites as appeared in Figure (1). Direct determination recorded high strength of electromagnetic fields in case of resistant mites to Vertimec on cotton followed by resistant strain on soybean with 45.37 and 28.14 G, respectively.0020 Therefore, there were specific differences in Axis Graph Magnetic Meter resulted of electromagnetic fields through both resistant and susceptible of adult females of *T.urticae* on certain crops as appeared in Figure (2).

Jonckheere-Terpstra Test showed that Std. Deviation of J-T Statistic=.957* among grouping variables of mites. Therefore, both Chi-Square and Median were =3.000a and 28.140^a. While, Kruskal-Wallis Test showed that Chi-Square=2.000^b at 5%. Proximity Matrix proved that a dissimilarity matrix with 51.587 as euclidean distance among variables while a similarity matrix was determined by the correlation between vectors of values was -.997.




Fig .1. Force of resulted electromagnetic fields through both resistant and susceptible of adult females of *T.urticae* on certain crops




Fig .2. Axis Graph Magnetic Meter resulted of electromagnetic fields through both resistant and susceptible of adult females of *T.urticae* on certain crops

Subsequently, toxicity of Vertimec was compared both of field and resistant strains with the susceptible strain of *T.urticae* on cotton. Whereas, LC₅₀ values were 1542.91, 1915.47 and 258.45 μLL⁻¹, respectively, which showed that LC₅₀ of cotton field strain exposed to moderately levels of CO₂ and UV recorded so close value to that of 20th generation of the laboratory resistant strain. The same situation was in the case of LC₉₀s which recorded 14207.39, 16328.37 and 1420.74 μLL⁻¹, respectively. Relative to the laboratory strain (S), the resistant ratios (RR) to Vertimec for *T.urticae* laboratory resistant strain and field strain, shown at Table (1). RR's at Vertimec LC₅₀s were 5.97- folds and 8.52- folds, respectively. The same trend occurred in case of assessment of Vertimec resistance in *T.urticae* strain on soybean.

Data showed that LC₅₀ values were 4013.25, 4220.17 and 198.82 μLL⁻¹, respectively, which showed that LC₅₀ of cotton field strain exposed to moderately levels of CO₂ and UV recorded a so close value to that of 20th generation of the laboratory resistant strain. The same situation was in the case of LC₉₀s which recorded 17210.74, 19102.35 and 780.34 μLL⁻¹, respectively. Relative to the laboratory strain (S), the resistant ratios (RR) to Vertimec for *T.urticae* laboratory resistant strain and field strain are shown at Table (1).

RR's at Vertimec LC₅₀'s were 5.97- folds and 8.52- folds, respectively. The same trend occurred in case of assessment of Vertimec resistance in *T.urticae* strain on soybean. Data showed that LC₅₀ values were 4013.25, 4220.17 and 198.82 μLL^{-1} , respectively, which showed that LC₅₀ of cotton field strain exposed to moderately levels of CO₂ and UV recorded a so close value to that of 20th generation of the laboratory resistant strain. The same situation was in the case of LC₉₀'s which recorded 17210.74, 19102.35 and 780.34 μLL^{-1} , respectively. Relative to the laboratory strain (S), the resistant ratios (RR) to

Vertimec for *T.urticae* laboratory resistant strain and field strain are shown at Table (2). RR's at Vertimec LC₅₀'s were 4.29- folds and 4.53- folds, respectively. According to Hayashi scale (1983), RRs of both strains resistant to Vertimec LC₅₀, infested cotton and soybean, could be ranked as low resistance. Friedman Test proved the highest significant effect of certain crops on LC₅₀'s values at 1% (Chi-Square =12.684**). Kendall's Coefficient of Concordance (Kendall's W^a) = .705** with highly significant Chi-Square =12.684 at 1%.

Table 1. Assessment of Vertimec toxicity against certain strains of adult females of *T.urticae*

<i>T.urticae</i> Strains	Crops	Vertimec Lethal Concentrations				LC ₉₀ /LC ₅₀ Ratio
		LC ₅₀		LC ₉₀		
		Main μLL^{-1}	*RR ₅₀	Main μLL^{-1}	*RR ₉₀	
Field	Cotton	1542.91 ^a	5.97	14207.39 ^a	9.99	9.21
Resistant		1915.47 ^a	8.52	16328.37 ^a	11.49	8.52
Susceptible		258.45 ^b		1420.74 ^b		5.50
Field	Soybean	4013.25 ^a	4.29	17210.74 ^a	22.10	4.29
Resistant		4220.17 ^a	4.53	19102.35 ^a	24.48	4.53
Susceptible		198.82 ^b		780.34 ^b		3.92

*Resistance ratio (RR) =LC₅₀ or LC₉₀ of Resistant Strain/ LC₅₀ or LC₉₀ of Susceptible Strain.

Pesticides resistance in *Tetranychus urticae* is a portent which caused by many reasons. One of their causes is the introduction of exceeding levels of UV and CO₂, which could assume a critical part to get a resistant field strain. Vertimec, an articulated miticide, LC₅₀ of the research center safe strain which is kept up under selection pressure till F40 and the resistant field strain in comparison with susceptible strain were 2099.38, 200.01 and 50.822 μLL^{-1} , respectively. Additionally, the raised esterases and mixed function oxidases (MFO) in both the laboratory and the field resistant strains had basically characterized the impact of extensive radiation of UV on the surpassed resistance levels recorded for the two strains (Abd El-Wahab and Taha, 2014).

Consequently, expanded CO₂ and global warming can be relied upon to positively influence the concoction barrier flagging framework in plants. Those factors will render them more vulnerable pest assault. The expanded number of produced generations every year and incessant populace flare-ups of potential pests require ceaseless uses of high measure of pesticides. It will boost exposed mites and insects to create pesticides' resistance vastly (Petzoldt and Seaman, 2007). Further, prolongation of insects' lifespan is prolonged under high CO₂ and temperature. Besides, such climate variations will stabilize such insecticide resistant in assortments of pests in their populations. Therefore, all mentioned will make more prominent harm plants even under broad pesticides measures. Also, a few classes of

pesticides have been appeared to be less powerful in controlling pests at higher temperatures (Musser and Shelton, 2005).

Vertimec (Abamectin) resistance in *T. urticae* was also reported by several authors (Beers et al., 1998). Stumpf and Nauen (2002) found that MFO (cytochrome P450-dependent monooxygenase) activity was higher in many strains of *T. urticae* than the susceptible strain. Consequently, Abamectin resistance was strongly synergized by PBO (piperonyl butoxide) and DEM (diethyl maleate), suggesting that MFO and GST (glutathione S-transferases) might be involved in abamectin resistance. Astonishingly, levels of cytochrome P450 monooxygenases, specifically, CYP6CM1 in mix with a simple to utilize counter acting agent identification framework permit a quick, dependable, and extremely touchy detection of pesticides' resistance to insects, and certainly to *Bemisia tabaci*, to Neonicotinoid and Pymetrozine (Nauen et al., 2013). Nevertheless, Li et al. (2016) demonstrated that RRs were ranged between 6.51 and 6.03 for *Myzus persicae* (Sulzer) infested tobacco, to certain pesticides in China. Results demonstrated that resistance ratios were RR = 6.51 and 6.03 which meant populaces have created minor imperviousness to Imidacloprid. One populace (NC) has achieved a high resistance level to Cyhalothrin (RR = 41.28), five populaces indicated medium level ($10.36 \leq RR \leq 20.45$), and the other six stayed powerless ($0.39 \leq RR \leq 3.53$). As respects Carbosulfan, three populaces have created medium resistance, four populaces indicated just minor resistance, and the other five ($0.81 \leq RR \leq 3.97$) were as yet vulnerable. Populace SZ built up a medium level (RR = 14.83) to Phoxim, the other 11 were vulnerable ($0.29 \leq RR \leq 2.41$). To examine the potential resistance system, restraint impacts of synergists and detoxifying compound exercises were identified. The outcomes showed that the MFO was the most vital detoxifying catalyst presenting Imidacloprid resistance, and CarE was most imperative to Cyhalothrin, Carbosulfan and Phoxim. Our examination gave an exhaustive overview of insecticide resistance of *M. persicae* in Chongqing, and proposed that distinctive districts should take comparing administration to postpone the insecticide resistance advancement and draw out the convenience of insecticides.

Furthermore, magnetic fields were tested on mites, *Tetranychus urticae* and *Polyphagotarsonemus latus*

latus which infested tomato leaves and some leaves were passed through 500 Gauss magnetic field and others were sprayed with magnetic water (Al-Ani, 2010). Results showed the significant decrease of mites' individuals and raised numbers of mites' eggs. That was explained upon hyperactivation of some enzymes in exposed mites to lay down more eggs. Even though, magnetic water showed its effect on the preparation of spray liquids against on the viability of plant protection agents as shown by Wachowiak and Kierzek (2002). They reported raised effectiveness in the control of *Phytophthora sp.* infested potato, after the use of these fungicides diluted in magnetic water. Likewise, magnetic water incremented the efficacy of acaricides against *Tetranychus urticae*. They were Talstar, Omite, Magus and Omite by the use of one and three magnetizers, one magnetizer and two semi-rings (Górski et al., 2009).

In addition, development of exposure methods to certain magnetic field is developed at this paper. Electromagnetic field as shown could play an important role to detect resistance of pesticides in mites through an electromagnetic interference (EMI) detector by translating electrophysiological changes that occurred mainly at glutamate-gated chloride channels (GluCl_s) and γ -aminobutyric acid-gated rdl and glutamate-gated GluCl_a chloride channels (GABACl) and resulted to Vertimec' resistance in mites (Riga et al., 2017). Even though and according to other studies which suggested detoxification enzymes with no effective target site were able to cause abamectin resistance in field *T. urticae* populations (Çağatay et al., 2018).

Near future, all recorded readings would be collected and stored at the cloud to be available whenever needed through Internet of Things system (IOT). Besides, data base was connected and joined with resistance ratios (RRs) which were gained from both fields and laboratory' data. Such system will be easier to be used by farmers wherever they are being. But utilizing the electromagnetic fields depending on physiological differences in resistant mites is appeared to be newly emergence trial through this paper. Besides, the solution of metabolic resistance in mites is already done by exposure to specific colors of light emitting diodes (LEDs). Each color could be linked to reduce metabolic pesticides resistance in specific pest on certain plant (Abd El-Wahab, 2015; Abd El-Wahab and Abouhatab, 2014; Abd El-Wahab and Bursic, 2014; Abd El-Wahab et

al.2014). Hence, LEDs with different colors are able to be connected with the full innovated system to do the two required steps: firstly, to detect the pesticides' resistance and secondly to stop or reduce it effectively.

IV. CONCLUSION

To conclude, as the pesticides' resistance is a big problem that leads to disasters affecting the environment, there is a new innovated instrument to detect and then reduce it. The solution is depending on differences in electromagnetic forces in resistant strains of *T.urticae* in comparison with susceptible strain. Then, resistance ratios have confirmed the presence of formed resistance. As a consequence, such device is capable to detect resistance to pesticides and farmers can count on it efficiently.

REFERENCES

- [1] R.A. Abd El-Wahab. "Direct effects of light emitting diodes (LEDs) on the two-spotted spider mite, *Tetranychus urticae*." Int. J. Sci. Res. Agricul. Sci., vol. 2, pp.79-85, 2015.
- [2] R.A. Abd El-Wahab. and E.E. Abouhatab. "Effects of light emitting diodes (LEDs) on the insect predators behavior against the two forms of *Tetranychus urticae*." Int. J. Chem. Biol. Sci. (IJCBS), vol. 1, no. 4, pp. 36-45, 2014.
- [3] R.A. Abd El-Wahab and V. Bursic. "Light emitting diodes (LEDs) reduce Vertimec, resistance in *Tetranychus urticae* (Koch)." Int. J. Chem. Biol. Sci. (IJCBS), vol. 1, no. 3, pp. 28-40, 2014.
- [4] R.A. Abd El-Wahab and T.M. Taha. "The relation between Vertimec resistance in the two-spotted spider mite, *Tetranychus urticae* and climate changes in Egypt." Int. J. Chem. Biol. Sci., vol. 1, no. 3pp. 1-10, 2014.
- [5] R.A. Abd El-Wahab, S. Lazic and V. Bursic. "Compatibility among insect predators and light emitting diodes (LEDs) against the two forms of *Tetranychus urticae* in greenhouses." Int. J. Chem. Biol. Sci. (IJCBS), vol. 1, no. 5, pp. 20-27, 2014.
- [6] W.S. Abbott. "A method of computing effectiveness of an insecticides." J. Econ. Entomol., vol 18, pp.265-267, 1925.
- [7] N. Al-Ani, "Effect of magnetic field on mites." J Al-Nahrain Univ., vol.13, no.3), pp.104-109, 2010.
- [8] E.H.Beers, H. Riedl and J.E.Dunley. "Resistance to abamectin and reversion to susceptibility to fenbutatin oxide in spider mite (Acari: Tetranychidae) populations in the Pacific Northwest." J. Econ. Entomol., vol. 91, pp. 352-360, 1998.
- [9] N.S. Çağatay, P. Menault, M. Riga, J. Vontas and A. Recep. "Identification and characterization of abamectin resistance," in *Tetranychus urticae*, Koch populations from greenhouses in Turkey, Crop Protection, vol. 112, pp.112–117, 2018.
- [10] V.V. Dawkar, Y.R. Chikate, P.R. Lomate, B.B. Dholakia, V.S. Gupta and A.P. Giri. "Molecular insights into resistance mechanisms of Lepidoptera insect pests against toxins." J. Proteome Res., vol. 12, no. 11, pp. 4727-4737, 2013.
- [11] V. Dittrich. "A comparative study of toxicological test methods on a population of the two-spotted spider mite, *T.urticae*." J. Econ. Entomol., vol 55, pp. 633-648, 1962.
- [12] G.P. Georgiou "Overview of insecticide resistance," in Managing Resistance to Agrochemicals Eds. MB Green, HM LeBaron, WK Moberg. pp. 18-41, American Chemical Society, 1990.
- [13] R. Gerharz. "Pest dislodgement by electromagnetic fields". <http://www.freepatentsonline.com/H000998.pdf> ,1991.
- [14] R. Górski , M. Wachowiak and M. Tomczak. "The effect of water magnetized with negative magnetic field on effectiveness of selected zoocides in the control of two-spotted spider mite (*Tetranychus urticae* Koch)." J. Plant Prot. Res., vol. 49, no. 1, pp. 87-91,2009.
- [15] A. Hayashi. "History, present status and management of insecticide resistance" in Pest Resistance to Pesticides. Eds J Fukami, Y Uesugi, K Ishizuka. pp. 31-53. Tokyo, Japan, Soft Science (Wash. D.C.), 1983.

- [16] K.R. Kranthi, Insecticide resistance -monitoring, mechanisms and management manual, Nagpur, India: CICR and ICAC, Washington. 2005, Pp.153.
- [17] LeOra Software. POLO-PC. A User's Guide to Probit or Logit Analysis LeOra Software, pp. 28, Berkeley, CA., 1994.
- [18] Y. Li, , Z. Xu, L. Shi, L. He and G. Shen. . "Insecticide resistance monitoring and metabolic mechanism study of the green peach aphid, *Myzus persicae* (Sulzer) (Hemiptera: Aphididae),in Chongqing, China." Pesticide Biochem.Physiol, vol.132, pp.21-28, 2016.
- [19] F.P. Musser and A.M. Shelton. "The influence of post-exposure temperature on the toxicity of insecticides to *Ostrinia nubilalis* (Lepidoptera:Crambidae)." Pest Manag Sci. vol. 61, pp. 508-510, 2005.
- [20] N.Nahar, W. Islam and M.M. Hague."Predation of three predators on two-spotted spider mite, *Tetranychus urticae* Koch (Acari: Tetranychidae)." J Life Earth Sci. vol. 1, pp. 1-4,2005.
- [21] R. Nauen, K. Raming and K.Wölfel. Online: 2013. Available: <http://www.google.com/patents/WO2013182613A1?cl=en>.
- [22] C. Petzoldt and A. Seaman. "Climate Change Effects on Insects and Pathogens. Fact Sheet." Online: 2007. Available: <http://www.climateandfarming.org/clrcc.php>.
- [23] M. Riga, S. Bajda, C. Themistokleous, S. PapaDaki, M. Palzewicz, W. Dermauw, J. Vontas and T. Van Leeuwen. "The relative contribution of target-site mutations in complex acaricide resistant phenotypes as assessed by marker assisted backcrossing in *Tetranychus urticae*." Scientific Reportss, vol. 7, pp. 9202-9214, 2017, DOI:10.1038/s41598-017-09054-y
- [24] N.Stumpf and R. Nauen. "Biochemical markers linked to abamectin resistance in *Tetranychus urticae* (Acari-Tetranychidae)." Pestic. Biochem. Physiol., vol. 72, pp. 111-121, 2002.
- [25] M.Wachowiak and R. Kierzek. "Wpływ dawki środka ochrony roślin, dodatku adiuwanta i wielkości kropeł na efektywność zwalczania agrofagów." Prog. Plant Protection/Post. Ochr.Roślin, vol. 43, no. 2, pp. 994–997, 2003.
- [26] X. Yang, L.L.K. Buschmann, Y. Zhu and D.C. Margolies" Susceptibility and detoxifying enzyme activity in two spider mite species (Acari : Tetranychidae) after selection with three secticides." J. Econ. Entomol., vol. 95, pp.399-406, 2002.